

**Unique Plan Description: ED Suspected Stroke VC**

**Plan Selection Display: ED Suspected Stroke VC**

**PlanType: Medical**

**Version: 1**

**Begin Effective Date: 10/30/2013 16:09**

**End Effective Date: Current**

**Available at all facilities**

**ED Suspected Stroke VC**

**Patient Care**

- Cardiac Monitoring  
*T;N, Constant Order, Stat, While in ED; Continuous if age >35*
- Communication Order  
*T;N, Activate Stroke Team*
- Neurological Checks  
*T;N, q1hr, Stat*
- Oxygen Therapy  
*T;N, Stat, Titrate O2 to maintain SpO2 of 94% or greater*
- Peripheral IV Insertion  
*T;N*
- Pulse Oximetry  
*T;N, Constant order, Stat, While in ED*
- Quality Measures, Stroke  
*T;N, Stat*
- Vital Signs  
*T;N, q15min, 1, hr*
- +1 Hours** Vital Signs  
*T;N, q30min, 1, hr*
- +2 Hours** Vital Signs  
*T;N, q1hr, Constant order*
- Access Portacath  
*T;N, PRN*
- Peripheral IV Insertion  
*T;N, PRN; second IV insertion. Use large bore catheter*

**Continuous Infusions**

- Sodium Chloride 0.9%  
*IV, Order Duration: 1 hr*

**Medications**

\*\*\*Refer to written tPa order set if tPa is indicated\*\*\*(NOTE)\*

- NS Bolus  
*500 mL, IV, Once, Max Bolus (mL/kg): 20, Order Duration: 1 doses (DEF)\*  
1,000 mL, IV, Once, Max Bolus (mL/kg): 20, Order Duration: 1 doses*
- labetalol  
*10 mg, IV Push, Once, Form: Injection (DEF)\*  
Comments: infuse over 1-2 minutes, may repeat x1, hold if heart rate less than 60  
20 mg, IV Push, Once, Form: Injection  
Comments: infuse over 1-2 minutes*
- nicardipine/NS 40 mg/200 mL Premix (IVS)\*  
Premix NS  
*Total Volume (mL): 200, IV  
niCARdipine IV additive  
40 mg, EB, 5 mg/hr*

**Laboratory**

- Beta hCG Qualitative Urine

*Urine, Stat collect, T;N*

*Comments: Females between 12-50 unless known hysterectomy*

- CBC w/ Differential  
*Blood, Stat Collect, T;N*
- CMP  
*Blood, Stat Collect, T;N*
- Istat Creatinine  
*Blood, Stat Collect, T;N*
- ISTAT MP  
*Stat Collect, T;N*
- PT/INR  
*Blood, Stat Collect, T;N*
- PTT  
*Blood, Stat Collect, T;N*
- Troponin  
*Blood, Stat Collect, T;N*
- Urinalysis with Culture if Indicated  
*Urine, Stat collect, T;N*

#### **Diagnostic Tests**

- CT Head or Brain w/o Contrast  
*T;N, Stat, Reason: Other (please specify), Reason: Stroke, Transport Mode: Cart*
- EKG  
*T;N, Stat, Other-Please Specify, Stroke*
- XR Chest 1 View Frontal  
*T;N, Stat, Reason: Other (please specify), Reason: Stroke, Transport Mode: Portable*
- CT Angiography Head/Neck w/+w/o Cont  
*T;N, Stat, Reason: Stroke, Transport Mode: Cart*  
*Comments: Creatinine 1.3- 1.7 use Visipaque (radi initiates); Creatinine > 1.7; Creatinine > 1.7 notify stroke provider prior to initiation notify stroke physician prior to initiation; allergy to iodine: find out reaction and notify stroke provider prior to initiation.*
- XR Chest 2 Views  
*T;N, Stat, Reason: Other (please specify), Reason: Stroke, Transport Mode: Cart*

#### **\*Report Legend:**

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase