

Unique Plan Description: MED Long-Term Central Catheter Insertion VC

Plan Selection Display: MED Long-Term Central Catheter Insertion VC

Plan Synonyms: Long-Term Central Catheter Insertion VC

PlanType: Medical

Version: 1

Begin Effective Date: 8/20/2013 12:40

End Effective Date: Current

Available at: **Ascension Health**
VC Rehab Hosp
VCH Behav Hlth
VCH HH Hospice
VCH Select
VCH St Francis
VCH St Joseph
VCH St Teresa
VCHSJ Assess
VCHSJ SBH
VCHST IP Rehab

MED Long-Term Central Catheter Insertion VC

Patient Care

- Obtain consent
T;N, For Central Line Catheter insertion. Call for procedural physician.
- Vital Signs
T;N, Per sedation protocol.
- Saline Lock Insert
T;N, Insert IV lock, discontinue prior to discharge for outpatients
- Central Line Care
T;N, qWeek, Per Protocol

Diet/Nutrition

- NPO
T;N

Medications

Give vancomycin if patient has cephalosporin allergy or severe penicillin allergy or at high risk for MRSA (chronic dialysis or MRSA colonization)**(NOTE)***

Ancef dosing based on weight as follows: - patient weight < 80 kg: give 1 gm - patient weight >= 80 kg: give 2 gm**(NOTE)***

- ceFAZolin
*1 gm, IV Piggyback, Once, Prophylaxis of infection: Pre-op, Form: Soln-IV (DEF)**
2 gm, IV Piggyback, Once, Prophylaxis of infection: Pre-op, Form: Soln-IV
- ***Vancomycin dosing based on weight as follows: - patient weight < 100 kg: give 1 gm - patient weight >= 100 kg: give 1.5 gm*****(NOTE)***
- vancomycin
*1 g, IV Piggyback, Once, Form: Soln-IV (DEF)**
1.5 g, IV Piggyback, Once, Form: Soln-IV
- Heparin/NS 2000 units/1000 mL (IVS)*
NS
Total Volume (mL): 1,000, Irrigation
Comments: apply on the sterile field for catheter preparation
heparin IV additive
2,000 units, EB
- MED Standard Flush VC(SUB)*
- lidocaine 1%/sodium bicarb 8.4% 20mL
1 syringes, IntraDermal, As Indicated, PRN Pain
Comments: to prevent pain from line insertion
- lidocaine 1% injectable solution

0.1 mL, IntraDermal, As Indicated, PRN Pain

Comments: to prevent pain from line insertion

- Xylocaine HCl with EPINEPHrine 1%-1:100,000 injectable solution
2 mL, IntraDermal, Form: Vial, Once, PRN Pain
Comments: administer up to 2 mL for local anesthesia
- Versed
2 mg, IV Push, Form: Injection, Once (DEF)*
1 mg, IV Push, Form: Injection, Once
0.5 mg, IV Push, Form: Injection, Once
- fentaNYL
50 mcg, IV Push, Form: Injection, Once (DEF)*
25 mcg, IV Push, Form: Injection, Once
75 mcg, IV Push, Form: Injection, Once
100 mcg, IV Push, Form: Injection, Once

Diagnostic Tests

- Chest XR 2 Views
T;N, Routine, Reason: Line Placement

Respiratory

- Oxygen Therapy
T;N, PRN

Communication Orders

- Communication Order
T;N, Have fluroscopy and ultrasound available

***Report Legend:**

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase